

Why artificial light at night should be a focus for global change research in the 21st century

Davies, Thomas W.; Smyth, Tim

Global Change Biology

DOI:
[10.1111/gcb.13927](https://doi.org/10.1111/gcb.13927)

Published: 01/03/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Davies, T. W., & Smyth, T. (2018). Why artificial light at night should be a focus for global change research in the 21st century. *Global Change Biology*, 24(3), 872-882.
<https://doi.org/10.1111/gcb.13927>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Why artificial light at night should be a focus for global change research in the
2 21st century

3 **Running head:** Artificial light is a global change issue

4 Thomas W. Davies^{1,2} and Tim Smyth³

5 ¹School of Ocean Sciences, Bangor University, Menai Bridge, UK

6 ²Centre for Geography, Environment and Society, University of Exeter, Penryn, UK

7 ³Plymouth Marine Laboratory, Prospect Place, Plymouth, Devon, UK

8 **Corresponding author:** Thomas Davies; School of Ocean Sciences, Bangor University, Menai
9 Bridge, UK; Thomas.Davies@exeter.ac.uk

10 **Key Words:** Artificial light at night, Global change, Ecology, Human health, Human-
11 environment interrelationships.

12 **Opinion**

13 Abstract

14 The environmental impacts of artificial light at night have been a rapidly growing field of global
15 change science in recent years. Yet light pollution has not achieved parity with other global change
16 phenomena in the level of concern and interest it receives from the scientific community, government
17 and non-governmental organisations. This is despite the globally widespread, expanding and changing
18 nature of night-time lighting; and the immediacy, severity and phylogenetic breadth of its impacts. In
19 this opinion piece, we evidence ten reasons why artificial light at night should be a focus for global
20 change research in the 21st century. Our reasons extend beyond those concerned principally with the
21 environment, to also include impacts on human health, culture and biodiversity conservation more
22 generally. We conclude that the growing use of night-time lighting will continue to raise numerous
23 ecological, human health and cultural issues, but that opportunities exist to mitigate its impacts by
24 combining novel technologies with sound scientific evidence. The potential gains from appropriate
25 management extend far beyond those for the environment, indeed it may play a key role in
26 transitioning towards a more sustainable society.

27 **Introduction**

28 While artificial light at night (ALAN) has been a long established man-made disturbance (Longcore
29 and Rich, 2004), the number of studies documenting its ecological and human health impacts has
30 grown dramatically in the last decade (Figure 1). Collectively, this body of research now highlights
31 the pervasiveness of ALAN’s impacts across a broad array of biomes, ecosystems, species and
32 behaviours. The measured biological responses occur at intensities and spectra of artificial light that
33 are currently encountered in the environment, and the global distribution of night-time lighting means
34 that it is likely already having widespread impacts in marine, freshwater and terrestrial habitats around
35 the world.

36

37 While ALAN research has gained notable momentum in recent years it is yet to achieve notoriety
38 among environmental scientists as a driver of global change. Here, we argue that ALAN should be a
39 focus for global change research in the 21st century. Our argument is broken down into ten points that
40 highlight the global extent of ALAN; the geographic scale of its influence; the potential to reverse its
41 environmental impacts; the rise of new human-environment conflicts with emerging lighting
42 technologies; its evolutionary novelty; the diverse array of species now known to be affected; the
43 extreme sensitivity of organisms to light; impacts on human health; cultural impacts on human-
44 environment interrelationships; and the feasibility of solutions. While we do not assert that ALAN is
45 any more important than other global change phenomena, we draw comparisons where they help
46 highlight the need for greater parity of concern.

47

48 **Globally widespread**

49 As with greenhouse gas emissions, ALAN is a globally widespread environmental pollutant. It is
50 estimated that 23% of the land surface between 75°N and 60°S (Falchi *et al.*, 2016), is exposed to
51 artificial skyglow (artificial light that is scattered in the atmosphere and reflected back to the ground).
52 This is comparable to the area of global ice-free land converted to either pasture or cropland,
53 estimated to be 35% in the year 2000 (Klein Goldewijk *et al.*, 2011). The degree of exposure

increased in all global terrestrial ecosystems between 2008 and 2012, with those important for biodiversity conservation often most affected (Bennie *et al.*, 2015b). Exposure to ALAN is not limited to terrestrial environments, with current best estimates indicating that 22% of the worlds' coastal regions (Davies *et al.*, 2014) are experiencing some degree of artificial illumination and 20% of marine protected areas are exposed across their entire range (Davies *et al.*, 2016). The amount of artificial light is also increasing in 13,061 terrestrial protected areas across Europe, Asia and South and Central America (Gaston *et al.*, 2015a), and 1,687 (14.7%) of the worlds marine protected areas (Davies *et al.*, 2016). Given that more than 95% of global population increases are projected to occur in the cities of economically developing countries over the next 50 years (Grimm *et al.*, 2008), and levels of light pollution are closely associated with population density and economic activity (Gallaway *et al.*, 2010), ALAN will continue to expand both in spatial extent and intensity throughout the 21st century without intervention.

Sphere of influence

Artificial light arises from point sources (municipal, industrial, commercial and residential), giving the impression that its impacts on the environment are highly localised. Indeed the majority of studies into the ecological impacts of ALAN quantify responses to direct lighting (Gaston *et al.*, 2015b). Artificial sky-glow increases the sphere of ALAN's potential influence far beyond a patch of habitat in the vicinity of a street light (Kyba & Hölker, 2013, Falchi *et al.*, 2016). Numerous taxa are adapted to make use of spatial and temporal patterns of natural sky brightness at intensities equivalent to or less than those created by artificial sky-glow (Naylor, 1999, Moore *et al.*, 2000, Dacke *et al.*, 2013, Last *et al.*, 2016, Warrant & Dacke, 2016), suggesting that lights in urban centres will have impacts in environments tens to hundreds of kilometres away. A dung beetle navigating its landscape using the Milky Way could, for example, become disorientated by artificial skyglow from a city tens or perhaps even hundreds of kilometres away (Kyba *et al.*, 2013), an effect comparable to a moth becoming disorientated by a street light hundreds of metres away (van Grunsven *et al.*, 2014).

81 While ALAN can be misconstrued as being a highly localised anthropogenic stressor, climate
82 warming is likewise misrepresented as globally widespread in its occurrence. Like ALAN,
83 ecologically relevant warming occurs at more localised spatial scales (Hannah *et al.*, 2014) (Figure 2),
84 and is influenced by variable topographical features such as slope and aspect that create refuges where
85 rates of warming are reduced (Bennie *et al.*, 2008, Maclean *et al.*, 2016). The ecological impacts of
86 climate change - like light pollution – are therefore likely to be spatially heterogeneous for organisms
87 with low mobility, but more widespread for taxa that depend on large scale movements for their
88 survival and reproduction. In the case of both stressors, population impacts on the former species are
89 manifest foremost through the loss and fragmentation of suitable habitat (Hannah *et al.*, 2014), while
90 impacts on the latter species are manifest via direct effects on population demography (Gaston &
91 Bennie, 2014).

92
93 **Lag effects**

94 Abating future rises in global temperatures constitutes one of the most significant challenges facing
95 humanity in the 21st century. Yet even if all fossil fuel combustion ceased with immediate effect, the
96 recovery of atmospheric CO₂ concentrations, global temperatures, ocean pH and oxygen
97 concentrations to pre-industrial levels would take hundreds to thousands of years (Frolicher *et al.*,
98 2014, Frölicher & Paynter, 2015, Mathesius *et al.*, 2015), and there is the very real possibility that
99 temperatures would continue to rise in the medium term (Frolicher *et al.*, 2014). By contrast, globally
100 widespread artificial light can be ‘switched off’ instantaneously. There would be no lag effect on the
101 physical environment following such an event, allowing the biological environment to immediately
102 begin the recovery process. While such a scenario would likely prove controversial, recent
103 technological advances present tangible ways of mitigating the ecological impacts of artificial light at
104 night (see reason ten). Failure to abate the environmental consequences of a man-made disturbance
105 using available viable solutions, would not inspire confidence in our ability to solve the apparently
106 insurmountable challenges posed by global climate change phenomena.

107

108 **The rise of LEDs**

109 Light-Emitting Diodes (LEDs) have grown from a 9% share of the lighting market in 2011 to 45% in
110 2014, and are forecast to reach 69% by 2020 (Zissis & Bertoldi, 2014). Their rising popularity stems
111 from the variety of colours that LEDs can be tailored to produce, their improved energy efficiency
112 over alternative electric light sources, and ability to produce ‘white’ light that is aesthetically pleasing
113 and provides enhanced visual performance (Schubert & Kim, 2005, Pimputkar *et al.*, 2009,). Whilst
114 LEDs are often advocated for their potential to reduce global CO₂ emissions, and the ability to tailor
115 their spectra to avoid unwanted environmental impacts (see ‘Feasibility of solutions’), environmental
116 scientists and human health experts have raised concerns about the broad-spectrum light (Davies *et*
117 *al.*, 2013, Macgregor *et al.*, 2014), and prominent short wavelength peak (Haim & Portnov, 2013,
118 Haim & Zubidat, 2015) that the commonly used white models emit (Figure 3).

119
120 Firstly, the broad range of wavelengths emitted by white LEDs likely enables organisms to perform
121 colour guided behaviours at night that were previously only possible during the day (Davies *et al.*,
122 2013). A range of intra and interspecific interactions could be affected including foraging (for
123 example seeking nectar rich flowers), predation (ability to locate and successfully capture prey),
124 sexual communication (ability to locate, identify and assess the fitness of conspecifics through visual
125 displays) and camouflage (ability to avoid detection by predators). Nocturnal species may find
126 themselves competing for resources with diurnal species where such interactions had previously not
127 existed (Macgregor *et al.*, 2014), and differences in the sensitivity of animal visual systems to white
128 LED light spectra could change the balance of species interactions (Davies *et al.*, 2013). Some
129 alternative lighting technologies also emit light across a broad range of wavelengths (for example
130 Metal Halide and Mercury Vapour lighting, Figure 3), however the energy efficiency of LEDs makes
131 them the lighting of choice in the 21st century, and as such research should focus on how any
132 unforeseen deleterious impacts can best be mitigated.

133

134 Secondly the short wavelength peak emitted by white LEDs coincides with the wavelengths to which
135 many biological responses are known to be sensitive. Many invertebrate behaviours (Cohen &
136 Forward, 2009, Gorbunov & Falkowski, 2002, Haddock *et al.*, 2010, van Langevelde *et al.*, 2011)
137 and the melatonin response (West *et al.*, 2011) are sensitive to short wavelengths of light (between
138 350 and 500nm), and some studies have demonstrated that white LED lighting has a greater impact on
139 short wavelength sensitive responses compared to alternative lighting technologies (Pawson & Bader,
140 2014).

141

142 Thirdly, because LEDs illuminate a broad range of wavelengths, they have the potential to affect a
143 greater variety of biological responses that are sensitive to specific wavelengths of light. To give one
144 example, while many invertebrate behaviours and the melatonin response are most sensitive to short
145 wavelength light, the phytochrome system in plants – which is associated with the timing of
146 flowering- is sensitive to red/far red light (660 and 720nm) (Bennie *et al.*, 2016). Using broad
147 wavelength light sources such as white LEDs therefore risks affecting more biological responses
148 across a greater variety of taxa than using narrow wavelength light sources such as low pressure
149 sodium lighting (Gaston *et al.*, 2012).

150

151 Fourthly, the improved energy efficiency offered by LEDs may encourage growth in the amount of
152 artificial light produced around the world. This ‘rebound effect’ can be observed in historical lighting
153 trends (see Kyba *et al.* 2014), and partly explains why aesthetic and decorative lighting installations
154 are now increasingly seen in municipal centres, on monuments, bridges and waterfront developments.

155

156 Finally, improvements in the energy efficiency of LED lighting coupled with the production
157 efficiency of solar cells is resulting in a rapid growth in off grid lighting installations, typically in
158 remote regions containing previously artificial light naive ecosystems (Mills & Jacobson 2007,
159 Adkins *et al.*, 2010, Dalberg Global Development Advisors 2013). The greatest ecological impacts of
160 ALAN over the next 50 years will likely occur in these previously artificial light naive regions, with
161 an ecology not previously shaped by night-time lighting.

162

163 **Evolutionary novelty**

164 Organisms have evolved with large scale fluctuations in atmospheric CO₂, climate temperatures and
165 ocean pH throughout history, while sudden changes to natural light regimes are unprecedented over
166 evolutionary time-scales. The harmonic movements of the earth, moon and sun provide reliable cues
167 to which many biological events are now highly attuned (Kronfeld-Schor *et al.*, 2013).

168 The ability of organisms to rapidly adapt to the introduction of ALAN through behavioural, genetic or
169 epigenetic changes is likely to be far more limited than for climate warming due to the unprecedented
170 nature of this change (Swaddle *et al.*, 2015). Further, the scattered growth of artificial lighting around
171 the world is a significant barrier to predicting where organisms will be able to seek out suitably dark
172 habitats in the future, and identifying where to allocate dark corridors that enable such migrations to
173 happen. While challenging, identifying where species need to go to survive climate warming, and how
174 they get there, is made simpler by the predictability of regional climatic shifts (for example poleward
175 migrations by land and sea, and upward migrations in high altitude regions) (Hannah *et al.*, 2007).

176

177 **Diversity of biological responses**

178 ALAN is now known to cause a plethora of environmental impacts from altering organism physiology
179 to changing the structure of ecological communities. The diversity of taxa affected continues to grow
180 and now includes birds (Kempnaers *et al.*, 2010, Dominoni, 2015), bats (Rydell, 1992, Stone *et al.*,
181 2009), sea turtles (Witherington, 1992, Kamrowski *et al.*, 2012), marsupials (Robert *et al.*, 2015),
182 rodents (Bird *et al.*, 2004), anurans (Hall, 2016); freshwater and marine fish (Becker *et al.*, 2012,
183 Riley *et al.*, 2013, Brüning *et al.*, 2015); moths (Frank, 1988, Wakefield *et al.*, 2015); beetles, spiders,
184 harvestmen, woodlice and ants (Davies *et al.*, 2012, Davies *et al.*, 2017); branchiopod (Moore *et al.*,
185 2000), amphipod (Davies *et al.*, 2012, Davies *et al.*, 2015, Navarro-Barranco & Hughes, 2015) and
186 copepod (Davies *et al.*, 2015) crustaceans; polychaete worms, colonial ascidians, and hydrozoans
187 (Davies *et al.*, 2015); corals (Kaniewska *et al.*, 2015), and terrestrial plants (Bennie *et al.*, 2015a,
188 Bennie *et al.*, 2016, French-Constant *et al.*, 2016). The documented impacts include those on animal

communication (Kempenaers *et al.*, 2010, van Geffen *et al.*, 2015), reproductive development (Dominoni *et al.*, 2013, Hansen *et al.*, 1992), the timing of reproduction (Kanievska *et al.*, 2015, Robert *et al.*, 2015), orientation (Frank, 1988, Witherington, 1992), habitat selection (Davies *et al.*, 2012, Davies *et al.*, 2015), predator avoidance (Wakefield *et al.*, 2015), predation pressure (Rydell, 1992, Becker *et al.*, 2012, Bolton *et al.*, 2017), circadian disruption (Brüning *et al.*, 2015, Raap *et al.*, 2015, Raap *et al.*, 2016), plant phenology (Bennie *et al.*, 2015a, Bennie *et al.*, 2016, ffrench-Constant *et al.*, 2016), and ecosystem services (Lewanzik & Voigt, 2014, Knop *et al.* 2017).

While those impacts on survival and reproductive success highlight that ALAN is likely causing widespread population losses for a variety of taxa, no population-level effects have so far been reliably demonstrated. This is in part because satellite images of night-time lights are not available in sufficiently high spatial resolution for inferences to be drawn regarding impacts on species populations that can be variable on the scale of tens to hundreds of metres (Elvidge *et al.* 2007). Disentangling the effects of street and residential lighting from those of urbanisation and land use change is challenging, since all of these explanatory variables likely contributes to population declines but all co-vary. Analyses using higher resolution images from the international space station (capable of identifying individual roads), may yield further insights, but tend to be focused on cities, preventing comparisons from being drawn across sufficiently large spatial scales. Recent developments in hemispherical photography allow ‘biologically relevant’ artificial skyglow to be mapped from ground level across thousands of square kilometres (Luginbuhl *et al.*, 2009, Zoltan, 2010), better enabling ecologists to quantify its impacts on populations of organisms that utilize celestial patterns of sky brightness, but perhaps not the population effects of direct lighting. Techniques to model the distribution of artificial light across towns and cities have also been developed (Bennie *et al.* 2014), however such models can be computationally expensive and have not yet been applied to the question of whether direct lighting has an impact on organism populations. Before After Control Impact (BACI) experiments have the potential to provide insights into the long-term responses of sessile species populations and those mobile taxa with <1km home ranges, however the finances and time required to implement them at appropriate spatial and temporal scales make this

approach less feasible in a limited funding environment. For now, quantifying the population level impacts of ALAN remains one of the most important and challenging problems facing ecologists working in this area.

Sensitivity of biological responses

Many organisms are extremely sensitive to natural light, utilizing light cues as dim as the Moon and the Milky Way to orientate themselves, navigate landscapes and identify conspecifics and resources at night (Ugolini *et al.*, 2005, Dacke *et al.*, 2013, Last *et al.*, 2016, Warrant & Dacke, 2016). Perhaps most striking is the growing number of documented responses to white LEDs in marine systems (Gorbunov & Falkowski, 2002, Davies *et al.*, 2015, Navarro-Barranco & Hughes, 2015, Bolton *et al.*, 2017), where species are both adapted to utilize short wavelengths that penetrate deeper in seawater, and are incredibly sensitive to natural light. Examples of this extreme sensitivity include copepods (*Calanus* sp.) that undergo diel vertical migration to depths of 50m guided only by variations in moonlight intensity during the arctic winter (Båtnes *et al.*, 2013, Last *et al.*, 2016); sessile invertebrate larvae that move and identify suitable settlement locations guided by light levels equivalent to moonless overcast nights (Thorson, 1964, Crisp & Ritz, 1973); and polychaete worms, corals and echinoderms that synchronise broadcast spawning events using monthly and annual variations in lunar light intensity (Naylor, 1999). Many of these responses are clearly sensitive enough to be affected both by direct lighting and artificial skyglow (Figure 4), and indeed such impacts have been demonstrated for zooplankton diel vertical migration in freshwater ecosystems (Moore *et al.*, 2000). Given the spatial extent of artificial skyglow in coastal regions (Davies *et al.*, 2014, Falchi *et al.*, 2016), the disproportionate importance of these regions for global biogeochemical cycles [coastal zones account for 30% of global ocean primary production but only 10% of global ocean surface area (Wollast, 1998)], and the role of diel vertical migration in maintaining these cycles (Hays, 2003), it is not unreasonable to surmise that ALAN could have detectable effects on ocean carbon and nutrient budgets in the near future.

Impacts on human health

In 2007, The World Health Organisation classified shift work that disrupted human circadian rhythms as a probable human carcinogen (International Agency for Research on Cancer, 2007). While this classification is primarily associated with shift work, exposure to ALAN has been linked to a variety of health disorders in people through the same circadian disruption mechanism. These include sleep disorders, depression, obesity, and the progression of some cancers (Cajochen *et al.*, 2011, Haim & Portnov, 2013, Chang *et al.*, 2014, Keshet-Sitton *et al.*, 2015). The prominent peak of blue wavelength light emitted by LEDs is of increasing concern, since it occurs at the most effective frequency for suppressing the production of melatonin (West *et al.*, 2011, Haim & Zubidat, 2015), a hormone released by the pineal gland that regulates sleep wake cycles and acts as an antioxidant. Over the last decade, LEDs have become a ubiquitous feature of human life, and can be found in street, residential, commercial and aesthetic lighting installations, laptops, televisions, e-readers, smart phones and tablets. Late evening exposure to LED light from handheld devices has been linked to circadian disruption of sleep wake cycles, and alertness and cognitive performance during the day (Cajochen *et al.*, 2011, Chang *et al.*, 2014).

The extent to which outdoor lighting impacts human health is yet to be reliably determined. While epidemiological studies have found correlations between the amount of outdoor lighting and some health effects (Kloog *et al.*, 2008, Koo *et al.*, 2016), as with ecological patterns they are limited by the inferences that can be drawn from satellite images (Defence Meteorological Satellite Programme Operational Line Scan) with insufficient spatial resolution (5km) to differentiate exposure to ALAN from other factors that co-vary across city districts at fine spatial scales (Elvidge *et al.* 2007, Kyba, 2016). The need for higher resolution images or novel approaches that can disentangle the effects on both ecology and human health of multiple urban pollutants that co-vary is clear, although individual level sensors can also reveal important impacts of daily light exposure on circadian disruption and stress (Figueiro *et al.* 2017). A more recent analysis using higher resolution (0.75km) images from the Visible Infrared Imaging Radiometer Suite (VIIRS) on board the Suomi National Polar-orbiting

Partnership satellite has revealed a significant association between ALAN and breast cancer incidence in the Greater Haifa Metropolitan Area in Israel (Rybnikova & Portnov, 2016). This analysis accounted for several potential co-varying explanatory factors, but not noise pollution, and atmospheric pollution explicitly.

Human-environment interrelationships

In a recent analysis that combined high resolution night-time satellite images with atmospheric dispersion models of artificial sky-glow, Falchi *et al.* (2016) estimated that more than 80% of the worlds' population currently live under light polluted skies, such that the Milky Way is hidden from one third of people alive today. This extraordinary change in our night-time environment escalated in the developed world during the mid to late 20th century, and is now rapidly transforming the cultures of billions in the developing world. The trend is concurrent with urbanisation [66% of the worlds' population will reside in urban areas by 2050 (United Nations, 2014)], and it contributes to the growing disconnect between people and nature that has become known as 'the extinction of experience' (Miller, 2005). This growing disconnect undermines public support for conservation issues by preventing individuals from connecting with, understanding, and forming attachments to the natural world (Miller, 2005).

The extinction of experience is another of the great challenges facing humanity in the 21st century. Miller (2005) argues it can be addressed by designing urban landscapes to facilitate 'meaningful interactions with the natural world'. There is perhaps no more profound way in which people can reconnect with nature, than giving them access to the Milky Way, and allowing them to experience the natural rhythms of moonlight and sunlight that they are evolutionarily pre-adapted to synchronise their physiology and behaviour with (Cajochen *et al.*, 2013, Wright Jr *et al.*, 2013). Like biodiversity conservation however, pristine skies have become tourist attractions restricted to regions awarded special status for their value to dark sky conservation (Collison & Poe, 2013, Rodrigues *et al.*, 2014, Pritchard, 2017) where many in the developed world can no longer afford to reside or visit. Pritchard

(2017) argues that dark sky protection programmes also risk suppressing the economic and cultural development of poorer nations in a way analogous to biodiversity conservation in the 20th century. In her appraisal of NASA's 'City Lights' composite satellite image of the world's lights at night (<http://earthobservatory.nasa.gov/Features/IntotheBlack/>) Pritchard (2017) warns against 'neo-colonial approaches to the conservation of natural night-sky brightness'. While it is clear the continued growth in artificial lighting risks perpetuating the disconnect between people and the environment - and this will inevitably contribute to the concomitant shifting baseline in conservation objectives (Pauly, 1995, Papworth *et al.*, 2009) – any intervention should seek to support the modernisation of societies while retaining their connections with the natural world. Pritchard (2017) describes achieving this balance as a 'new frontier in 21st century conservation'.

Feasibility of solutions

While the recent growth in LED lighting has raised concerns among environmental scientists and human health experts, this technology offers lighting managers greater flexibility when it comes to tailoring the timing, intensity and spectral power distribution of municipal lighting systems (Gaston, 2013, Davies *et al.*, 2017). Of the local authorities in England, 23% are engaged in permanent part-night lighting schemes where street lights are turned off between midnight and 04:00 to 05:00 AM, while 39% are engaged in permanent dimming schemes where lights are dimmed for at least some period of the night (Campaign to Protect Rural England, 2014). Increasing constraints on local authority budgets have incentivized the adoption of these lighting strategies in the wake of the 2008 global financial crash, however more often the reasons given for their implementation are improved energy savings and reduced CO₂ emissions. Both dimming and part-night lighting are better enabled by switching to LED, and introducing central management systems that use wireless communication technology to programme individual street lights remotely.

The ecological benefits of dimming and part-night lighting are not yet well explored (although see Azam *et al.*, 2015, Day *et al.*, 2015, Davies *et al.*, 2017). A recent emphasis in the ecological

literature has instead been on tailoring spectral power distributions to reduce known ecological impacts (Pawson & Bader, 2014, Longcore *et al.*, 2015, Brüning *et al.*, 2016, Rivas *et al.*, 2015, Spoelstra *et al.*, 2015, van Geffen *et al.*, 2015, Davies *et al.*, 2017), despite this approach being less popular among lighting managers and engineers who often focus on the improved visual performance offered by broad spectrum lighting as a key selling point. These studies collectively present an inconsistent picture of whether spectral manipulation can be used to effectively mitigate the ecological impacts of ALAN. This is partly because some studies compare narrow spectrum (for example red, green and blue) light with broad spectrum light sources, while others either decrease the amount of light occurring at wavelengths known to manifest certain ecological responses (usually shorter wavelengths in the visible spectrum), or increase the amount of light occurring at wavelengths that do not give rise to these responses (longer wavelengths in the visible spectrum). Even if a unified approach were adopted in spectral manipulation experiments, it seems unlikely that a publically acceptable light spectra that does not give rise to any ecological impacts can be developed, because different species responses are evolutionarily adapted to utilize different wavelengths of light.

Examples of this are abundant in the emerging literature on the ecological impacts of artificial light. The number of beetle taxa aggregating under white LED lighting can be reduced by switching to amber, but this has no discernible effect on the number of spider taxa that aggregate (Davies *et al.*, 2017). Many animal responses are sensitive to short wavelength light (van Langevelde *et al.*, 2011, Rivas *et al.*, 2015, Spoelstra *et al.*, 2017), while phenological responses in plants are most sensitive to the longer wavelengths recommended to avoid such effects (Bennie *et al.*, 2015a, Bennie *et al.*, 2016). Male caterpillars of the moth *Mamestra brassicae* reared under green artificial light reached a lower maximum mass, pupated earlier and obtained a lower pupal mass than those reared under red light (van Geffen *et al.* 2014), while red light inhibited the attractiveness of a female pheromone lure to more adult males of the winter moth *Operophtera brumata* than did green light (van Geffen *et al.* 2015).

Studies investigating the ecological benefits of part-night lighting have also highlighted that different taxa respond in different ways (Azam *et al.*, 2015, Day *et al.*, 2015, Davies *et al.*, 2017), and the adoption of part-night lighting schemes is often inhibited by a perception among political actors that they lack popular support. There are both perceived and realised benefits of artificial light for society, including in the areas of road safety, crime, and the economy (Gaston *et al.*, 2015c). The night time economy in the UK, for example, was worth £67bn in 2016 (MAKE Associates, 2017), and accounted for up to 27% of town and city centre turnover and 10% of most locations overall employment figure in 2009 (VisitEngland, 2012).

While modern lighting technologies offer the potential to reduce the ecological impacts of ALAN, identifying how this is best achieved is clearly complex. Studies are needed across a wide variety of taxonomic groups and lighting approaches, to develop options that are both socially and ecologically acceptable.

Conclusion

Research into the ecological, human health and societal consequences of ALAN is now growing rapidly. Here, we have highlighted ten reasons why ALAN should, and likely will be a focus for global change research in the 21st century. Most important to consider, is the notion that while ALAN is having widespread and profound impacts on people and the environment, strategies for abating them are already being explored. Solving the challenges posed by ALAN would not only improve environmental and human health outcomes, but also enhance the human experience of nature and change perceptions of the natural world in a way that facilitates the necessary transition towards a more environmentally orientated and hence sustainable society. It would also inspire greater confidence in our ability to tackle the problems posed by other global change phenomena. The challenge now is identifying how best to address to the complex array of ecological, human health and cultural problems presented by society's propensity for illuminating the night.

Acknowledgements

The authors would like to thank Dr Ilya Mclean (University of Exeter) for granting permissions to use images in Figure 2. The work leading to this publication was supported by a Welsh Government and European Regional Development Fund funded Sêr Cymru II Fellowship awarded to TWD.

References

- Adkins E, Eapen S, Kaluwile F, Nair G, Modi V (2010) Off-grid energy services for the poor: introducing LED lighting in the Millennium Villages Project in Malawi. *Energy Policy*, **38**, 1087-1097.
- Azam C, Kerbiriou C, Vernet A, Julien J-F, Bas Y, Plichard L,...Le Viol I (2015) Is part-night lighting an effective measure to limit the impacts of artificial lighting on bats? *Global Change Biology*, **21**, 4333-4341.
- Båtnes AS, Miljeteig C, Berge J, Greenacre M, Johnsen G (2013) Quantifying the light sensitivity of *Calanus* spp. during the polar night: potential for orchestrated migrations conducted by ambient light from the sun, moon, or aurora borealis? *Polar Biology*, **38**, 51-65.
- Becker A, Whitfield AK, Cowley PD, Järnegen J, Næsje TF (2012) Potential effects of artificial light associated with anthropogenic infrastructure on the abundance and foraging behaviour of estuary-associated fishes. *Journal of Applied Ecology*, **50**, 43-50.
- Bennie J, Davies TW, Cruse D, Gaston KJ (2016) Ecological effects of artificial light at night on wild plants. *Journal of Ecology*, **104**, 611-620.
- Bennie J, Davies TW, Inger R, Gaston KJ (2014) Mapping artificial lightscapes for ecological studies. *Methods in Ecology and Evolution*, **5**, 534-540.
- Bennie J, Davies TW, Cruse D, Inger R, Gaston KJ (2015a) Cascading effects of artificial light at night: resource-mediated control of herbivores in a grassland ecosystem. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **370**, 20140131.

- 405 Bennie J, Duffy J, Davies T, Correa-Cano M, Gaston K (2015b) Global trends in exposure to light
406 pollution in natural terrestrial ecosystems. *Remote Sensing*, **7**, 2715-2730.
- 407 Bennie J, Huntley B, Wiltshire A, Hill MO, Baxter R (2008) Slope, aspect and climate: spatially
408 explicit and implicit models of topographic microclimate in chalk grassland. *Ecological*
409 *Modelling*, **216**, 47-59.
- 410 Bird BL, Branch LC, Miller DL (2004) Effects of coastal lighting on foraging behavior of beach mice.
411 *Conservation Biology*, **18**, 1435-1439.
- 412 Bolton D, Mayer-Pinto M, Clark GF, Dafforn KA, Brassil WA, Becker A, Johnston EL (2017)
413 Coastal urban lighting has ecological consequences for multiple trophic levels under the sea.
414 *Science of the Total Environment*, **576**, 1-9.
- 415 Brüning A, Hölker F, Franke S, Kleiner W, Kloas W (2016) Impact of different colours of artificial
416 light at night on melatonin rhythm and gene expression of gonadotropins in European perch.
417 *Science of The Total Environment*, **543**, 214-222.
- 418 Brüning A, Hölker F, Franke S, Preuer T, Kloas W (2015) Spotlight on fish: light pollution affects
419 circadian rhythms of European perch but does not cause stress. *Science of the Total*
420 *Environment*, **511**, 516-522.
- 421 Cajochen C, Altanay-Ekici S, Münch M, Frey S, Knoblauch V, Wirz-Justice A (2013) Evidence that
422 the lunar cycle influences human sleep. *Current Biology*, **23**, 1485-1488.
- 423 Cajochen C, Frey S, Anders D, Späti J, Bues M, Pross A,...Stefani O (2011) Evening exposure to a
424 light-emitting diodes (LED)-backlit computer screen affects circadian physiology and
425 cognitive performance. *Journal of Applied Physiology*, **110**, 1432-1438.
- 426 Chang A-M, Aeschbach D, Duffy JF, Czeisler CA (2014) Evening use of light-emitting eReaders
427 negatively affects sleep, circadian timing, and next-morning alertness. *Proceedings of the*
428 *National Academy of Sciences of the United States of America*. **112**, 1232-1237.
- 429 Cohen JH, Forward RB (2009) Zooplankton diel vertical migration - a review of proximate control.
430 *Oceanography and Marine Biology: An Annual Review*, **47**, 77-110.
- 431 Collison FM, Poe K (2013) "Astronomical Tourism": The Astronomy and Dark Sky Program at Bryce
432 Canyon National Park. *Tourism Management Perspectives*, **7**, 1-15.

- 433 Campaign to Protect Rural England (2014) Shedding Light: A survey of local authority approaches to
434 lighting in England, London.
- 435 Crisp DJ, Ritz DA (1973) Responses of cirripede larvae to light. I. Experiments with white light.
436 Marine Biology, **23**, 327-335.
- 437 Dacke M, Baird E, Byrne M, Scholtz CH, Warrant EJ (2013) Dung beetles use the Milky Way for
438 orientation. Current Biology, **23**, 298-300.
- 439 Dalberg Global Development Advisors (2013) Lighting Africa Market Trends Report 2012. In:
440 *Lighting Africa*.
- 441 Davies TW, Bennie J, Cruse D, Blumgart D, Inger R, Gaston KJ (2017) Multiple night-time light-
442 emitting diode lighting strategies impact grassland invertebrate assemblages. Global Change
443 Biology, **23**, 2641-2648.
- 444 Davies TW, Bennie J, Gaston KJ (2012) Street lighting changes the composition of invertebrate
445 communities. Biology Letters, **8**, 764-767.
- 446 Davies TW, Bennie J, Inger R, De Ibarra NH, Gaston KJ (2013) Artificial light pollution: are shifting
447 spectral signatures changing the balance of species interactions? Global Change Biology, **19**,
448 1417-1423.
- 449 Davies TW, Coleman M, Griffith KM, Jenkins SR (2015) Night-time lighting alters the composition
450 of marine epifaunal communities. Biology Letters, **11**, 20150080.
- 451 Davies TW, Duffy JP, Bennie J, Gaston KJ (2014) The nature, extent, and ecological implications of
452 marine light pollution. Frontiers in Ecology and the Environment, **12**, 347-355.
- 453 Davies TW, Duffy JP, Bennie J, Gaston KJ (2016) Stemming the tide of light pollution encroaching
454 into marine protected areas. Conservation Letters, **9**, 164-171.
- 455 Day J, Baker J, Schofield H, Mathews F, Gaston KJ (2015) Part-night lighting: implications for bat
456 conservation. Animal Conservation, **18**, 512-516.
- 457 Dominoni D (2015) The effects of light pollution on biological rhythms of birds: an integrated,
458 mechanistic perspective. Journal of Ornithology, **156**, 409-418.
- 459 Dominoni D, Quetting M, Partecke J (2013) Artificial light at night advances avian reproductive
460 physiology. Proceedings of the Royal Society B: Biological Sciences, **280**, 20123017.

- 461 Elvidge CD, Cinzano P, Pettit D, Arvesen J, Sutton P, Small C,...Weeks J (2007) The Nightsat
462 mission concept. *International Journal of Remote Sensing*, **28**, 2645-2670.
- 463 Falchi F, Cinzano P, Duriscoe D, Kyba CC, Elvidge CD, Baugh K,...Furgoni R (2016) The new
464 world atlas of artificial night sky brightness. *Science Advances*, **2**, e1600377.
- 465 Ffrench-Constant RH, Somers-Yeates R, Bennie J, Economou T, Hodgson D, Spalding A, McGregor
466 PK (2016) Light pollution is associated with earlier tree budburst across the United Kingdom.
467 *Proceedings of the Royal Society B: Biological Sciences*, **283**, No 1833.
- 468 Figueiro MG, Steverson B, Heerwagen J, Kampschroer K, Hunter CM, Gonzales K,...Rea MS (2017)
469 The impact of daytime light exposures on sleep and mood in office workers. *Sleep Health*, **3**,
470 204-215.
- 471 Forward RB, Cronin TW, Stearns DE (1984) Control of diel vertical migration: Photoresponses of a
472 larval crustacean1. *Limnology and Oceanography*, **29**, 146-154.
- 473 Frank KD (1988) Impact of outdoor lighting on moths an assessment. *Journal of the Lepidopterists'*
474 *Society*, **42**, 63-93.
- 475 Frölicher TL, Paynter DJ (2015) Extending the relationship between global warming and cumulative
476 carbon emissions to multi-millennial timescales. *Environmental Research Letters*, **10**, 075002.
- 477 Frolicher TL, Winton M, Sarmiento JL (2014) Continued global warming after CO₂ emissions
478 stoppage. *Nature Climate Change*, **4**, 40-44.
- 479 Gallaway T, Olsen RN, Mitchell DM (2010) The economics of global light pollution. *Ecological*
480 *Economics*, **69**, 658-665.
- 481 Gaston KJ (2013) Sustainability: A green light for efficiency. *Nature*, **497**, 560-561.
- 482 Gaston KJ, Bennie J (2014) Demographic effects of artificial nighttime lighting on animal
483 populations. *Environmental Reviews*, **22**, 323-330.
- 484 Gaston KJ, Davies TW, Bennie J, Hopkins J (2012) Reducing the ecological consequences of night-
485 time light pollution: options and developments. *Journal of Applied Ecology*, **49**, 1256-1266.
- 486 Gaston KJ, Duffy JP, Bennie J (2015a) Quantifying the erosion of natural darkness in the global
487 protected area system. *Conservation Biology*, **29**, 1132-1141.

- 488 Gaston KJ, Gaston S, Bennie J, Hopkins J (2015c) Benefits and costs of artificial nighttime lighting of
489 the environment. *Environmental Reviews*, **23**, 14-23.
- 490 Gaston KJ, Visser ME, Hölker F (2015b) The biological impacts of artificial light at night: the
491 research challenge. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
492 **370**, 20140133.
- 493 Gorbunov MY, Falkowski PG (2002) Photoreceptors in the cnidarian hosts allow symbiotic corals to
494 sense blue moonlight. *Limnology and Oceanography*, **47**, 309-315.
- 495 Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM (2008) Global change
496 and the ecology of cities. *Science*, **319**, 756-760.
- 497 Haddock SHD, Moline MA, Case JF (2010) Bioluminescence in the sea. *Annual Review of Marine*
498 *Science*, **2**, 443-493.
- 499 Haim A, Portnov BA (2013) Light pollution as a new risk factor for human breast and prostate
500 cancers, New York, Springer.
- 501 Haim A, Zubidat AE (2015) LED light between Nobel Prize and cancer risk factor. *Chronobiology*
502 *International*, **32**, 725-727.
- 503 Hall AS (2016) Acute artificial light diminishes central Texas anuran calling behavior. *The American*
504 *Midland Naturalist*, **175**, 183-193.
- 505 Hannah L, Flint L, Syphard AD, Moritz MA, Buckley LB, Mccullough IM (2014) Fine-grain
506 modeling of species' response to climate change: holdouts, stepping-stones, and microrefugia.
507 *Trends in Ecology & Evolution*, **29**, 390-397.
- 508 Hannah L, Midgley G, Andelman S, Araújo M, Hughes G, Martinez-Meyer E,...Williams P (2007)
509 Protected area needs in a changing climate. *Frontiers in Ecology and the Environment*, **5**, 131-
510 138.
- 511 Hansen T, Stefansson S, Taranger GL (1992) Growth and sexual maturation in Atlantic salmon,
512 *Salmo salar* L., reared in sea cages at two different light regimes. *Aquaculture Research*, **23**,
513 275-280.
- 514 Hays GC (2003) A review of the adaptive significance and ecosystem consequences of zooplankton
515 diel vertical migrations. *Hydrobiologia*, **503**, 163-170.

- 516 International Agency for Research on Cancer (2007) IARC Monographs on the evaluation of
 517 carcinogenic risks to humans Volume 98: Painting, firefighting, and shiftwork. In: *World*
 518 *Heath Organisation International Agency for Research on Cancer*, Lyon 2010.
- 519 Kamrowski RL, Limpus C, Moloney J, Hamann M (2012) Coastal light pollution and marine turtles:
 520 assessing the magnitude of the problem. *Endangered Species Research*, **19**, 85-98.
- 521 Kaniewska P, Alon S, Karako-Lampert S, Hoegh-Guldberg O, Levy O (2015) Signaling cascades and
 522 the importance of moonlight in coral broadcast mass spawning. *eLife*, **4**, e09991.
- 523 Kempenaers B, Borgström P, Loës P, Schlicht E, Valcu M (2010) Artificial night lighting affects
 524 dawn song, extra-pair siring success, and lay date in songbirds. *Current Biology*, **20**, 1735-
 525 1739.
- 526 Keshet-Sitton A, Or-Chen K, Yitzhak S, Tzabary I, Haim A (2015) Can avoiding light at night reduce
 527 the risk of breast cancer? *Integrative Cancer Therapies*, **1534735415618787**.
- 528 Klein Goldewijk K, Beusen A, van Drecht G, De Vos M (2011) The HYDE 3.1 spatially explicit
 529 database of human-induced global land-use change over the past 12,000 years. *Global*
 530 *Ecology and Biogeography*, **20**, 73-86.
- 531 Kloog I, Haim A, Stevens RG, Barchana M, Portnov BA (2008) Light at night co-distributes with
 532 incident breast but not lung cancer in the female population of Israel. *Chronobiology*
 533 *International*, **25**, 65-81.
- 534 Knop E, Zoller L, Ryser R, Gerpe C, Hörler M, Fontaine C (2017) Artificial light at night as a new
 535 threat to pollination. *Nature*, **548**, 206-209.
- 536 Koo YS, Song J-Y, Joo E-Y, Lee H-J, Lee E, Lee S-K, Jung K-Y (2016) Outdoor artificial light at
 537 night, obesity, and sleep health: Cross-sectional analysis in the KoGES study. *Chronobiology*
 538 *International*, **33**, 301-314.
- 539 Kronfeld-Schor N, Dominoni D, De La Iglesia H, Levy O, Herzog ED, Dayan T, Helfrich-Forster C
 540 (2013) Chronobiology by moonlight. *Proceedings of the Royal Society B: Biological*
 541 *Sciences*, **280**, 20123088.
- 542 Kyba CM, Hölker F (2013) Do artificially illuminated skies affect biodiversity in nocturnal
 543 landscapes? *Landscape Ecology*, **28**, 1637-1640.

- 544 Kyba C, Hänel A, Hölker F (2014) Redefining efficiency for outdoor lighting. *Energy &*
545 *Environmental Science*, **7**, 1806-1809.
- 546 Kyba CCM (2016) Defense Meteorological Satellite Program data should no longer be used for
547 epidemiological studies. *Chronobiology International*, **33**, 943-945.
- 548 Last KS, Hobbs L, Berge J, Brierley AS, Cottier F (2016) Moonlight drives ocean-scale mass vertical
549 migration of zooplankton during the Arctic winter. *Current Biology*, **26**, 244-251
- 550 Lewanzik D, Voigt CC (2014) Artificial light puts ecosystem services of frugivorous bats at risk.
551 *Journal of Applied Ecology*, **51**, 388-394.
- 552 Longcore T, Aldern HL, Eggers JF, Flores S, Franco L, Hirshfield-Yamanishi E,...Barroso AM
553 (2015) Tuning the white light spectrum of light emitting diode lamps to reduce attraction of
554 nocturnal arthropods. *Philosophical Transactions of the Royal Society B*, **370**, 20140125.
- 555 Longcore T, Rich C (2004) Ecological light pollution. *Frontiers in Ecology and the Environment*, **2**,
556 191-198.
- 557 Luginbuhl CB, Duriscoe DM, Moore CW, Richman A, Lockwood GW, Davis DR (2009) From the
558 ground up II: sky glow and near-ground artificial light propagation in Flagstaff, Arizona.
559 *Publications of the Astronomical Society of the Pacific*, **121**, 204-212.
- 560 Macgregor CJ, Pocock MJO, Fox R, Evans DM (2014) Pollination by nocturnal Lepidoptera, and the
561 effects of light pollution: a review. *Ecological Entomology*, **40**, 187-198.
- 562 Maclean IMD, Suggitt AJ, Wilson RJ, Duffy JP, Bennie JJ (2016) Fine-scale climate change:
563 modelling spatial variation in biologically meaningful rates of warming. *Global Change*
564 *Biology*, **23**, 253-268.
- 565 MAKE Associates Ltd. (2017) The NightMix index.
- 566 Mathesius S, Hofmann M, Caldeira K, Schellnhuber HJ (2015) Long-term response of oceans to CO₂
567 removal from the atmosphere. *Nature Climate Change*, **5**, 1107-1113.
- 568 Miller JR (2005) Biodiversity conservation and the extinction of experience. *Trends in ecology &*
569 *evolution*, **20**, 430-434.

- 570 Mills E, Jacobson A (2007) The off-grid lighting market in western Kenya: LED alternatives and
571 consumer preferences in a Millennium Development village. Lumina Project Technical
572 Report.
- 573 Moore MV, Pierce SM, Walsh HM, Kvalvik SK, Lim JD (2000) Urban light pollution alters the diel
574 vertical migration of *Daphnia*. Verhandlungen des Internationalen Verein Limnologie, **27**,
575 779-782.
- 576 Navarro-Barranco C, Hughes LE (2015) Effects of light pollution on the emergent fauna of shallow
577 marine ecosystems: Amphipods as a case study. Marine Pollution Bulletin, **94**, 235-240.
- 578 Naylor E (1999) Marine animal behaviour in relation to lunar phase. Earth, Moon, and Planets, **85**,
579 291-302.
- 580 Papworth SK, Rist J, Coad L, Milner-Gulland EJ (2009) Evidence for shifting baseline syndrome in
581 conservation. Conservation Letters, **2**, 93-100.
- 582 Pauly D (1995) Anecdotes and the shifting baseline syndrome of fisheries. Trends in Ecology and
583 Evolution, **10**, 430.
- 584 Pawson SM, Bader MKF (2014) LED lighting increases the ecological impact of light pollution
585 irrespective of color temperature. Ecological Applications, **24**, 1561-1568.
- 586 Pimputkar S, Speck JS, Denbaars SP, Nakamura S (2009) Prospects for LED lighting. Nature
587 Photonics, **3**, 180-182.
- 588 Pritchard SB (2017) The trouble with darkness: NASA's Suomi satellite images of Earth at night.
589 Environmental History, **22**, 312-330.
- 590 Raap T, Pinxten R, Eens M (2015) Light pollution disrupts sleep in free-living animals. Scientific
591 Reports, **5**, 13557.
- 592 Raap T, Pinxten R, Eens M (2016) Artificial light at night disrupts sleep in female great tits (*Parus*
593 major) during the nestling period, and is followed by a sleep rebound. Environmental
594 Pollution, **215**, 125-134.
- 595 Riley WD, Davison PI, Maxwell DL, Bendall B (2013) Street lighting delays and disrupts the
596 dispersal of Atlantic salmon (*Salmo salar*) fry. Biological Conservation, **158**, 140-146.

- 597 Rivas ML, Tomillo PS, Uribeondo JD, Marco A (2015) Leatherback hatchling sea-finding in response
598 to artificial lighting: Interaction between wavelength and moonlight. *Journal of Experimental*
599 *Marine Biology and Ecology*, **463**, 143-149.
- 600 Robert KA, Lesku JA, Partecke J, Chambers B (2015) Artificial light at night desynchronizes strictly
601 seasonal reproduction in a wild mammal. *Proceedings of the Royal Society B: Biological*
602 *Sciences*, **282**, 20151745.
- 603 Rodrigues ALO, Rodrigues A, Peroff DM (2014) The sky and sustainable tourism development: a
604 case study of a dark sky reserve implementation in Alqueva. *International Journal of Tourism*
605 *Research*, **17**, 292-302.
- 606 Rybnikova NA, Portnov BA (2016) Outdoor light and breast cancer incidence: a comparative analysis
607 of DMSP and VIIRS-DNB satellite data. *International Journal of Remote Sensing*, 1-10.
- 608 Rydell J (1992) Exploitation of insects around streetlamps by bats in Sweden. *Functional Ecology*, **6**,
609 744-750.
- 610 Schubert EF, Kim JK (2005) Solid-state light sources getting smart. *Science*, **308**, 1274-1278.
- 611 Spoelstra K, van Grunsven RHA, Donners M, Gienapp P, Huigens ME, Slaterus R,... Veenendaal E
612 (2015) Experimental illumination of natural habitat: an experimental set-up to assess the
613 direct and indirect ecological consequences of artificial light of different spectral
614 composition. *Philosophical Transactions of the Royal Society B*, **370**, 20140129.
- 615 Spoelstra K, van Grunsven RHA, Ramakers JJC, Ferguson KB, Raap T, Donners M,... Visser ME
616 (2017) Response of bats to light with different spectra: light-shy and agile bat presence is
617 affected by white and green, but not red light. *Proceedings of the Royal Society B: Biological*
618 *Sciences*, **284**, 20170075.
- 619 Stone EL, Jones G, Harris S (2009) Street lighting disturbs commuting bats. *Current Biology*, **19**,
620 1123-1127.
- 621 Svane I, Dolmer P (1995) Perception of light at settlement: a comparative study of two invertebrate
622 larvae, a scyphozoan planula and a simple ascidian tadpole. *Journal of Experimental Marine*
623 *Biology and Ecology*, **187**, 51-61.

- 624 Swaddle JP, Francis CD, Barber JR, Cooper CB, Kyba CC, Dominoni DM,...Luther D (2015) A
 625 framework to assess evolutionary responses to anthropogenic light and sound. *Trends in*
 626 *Ecology & Evolution*, **30**, 550-560.
- 627 Tankersley RA, Mckelvey LM, Forward RB (1995) Responses of estuarine crab megalopae to
 628 pressure, salinity and light: Implications for flood-tide transport. *Marine Biology*, **122**, 391-
 629 400.
- 630 Thorson G (1964) Light as an ecological factor in the dispersal and settlement of larvae of marine
 631 bottom invertebrates. *Ophelia*, **1**, 167-208.
- 632 Ugolini A, Boddi V, Mercatelli L, Castellini C (2005) Moon orientation in adult and young
 633 sandhoppers under artificial light. *Proceedings of the Royal Society B: Biological Sciences*,
 634 **272**, 2189-2194.
- 635 United Nations Population Division (2014) World Urbanization Prospects: The 2014 Revision,
 636 Highlights (ST/ESA/SER.A/352).
- 637 van Geffen KG, van Grunsven RHA, van Ruijven J, Berendse F, Veenendaal EM (2014) Artificial
 638 light at night causes diapause inhibition and sex-specific life history changes in a moth.
 639 *Ecology and Evolution*, **4**, 2802-2809.
- 640 van Geffen KG, van Eck E, De Boer RA, van Grunsven RHA, Salis L, Berendse F, Veenendaal EM
 641 (2015) Artificial light at night inhibits mating in a Geometrid moth. *Insect Conservation and*
 642 *Diversity*, **8**, 282-287.
- 643 van Grunsven RHA, Lham D, van Geffen KG, Veenendaal EM (2014) Range of attraction of a 6-W
 644 moth light trap. *Entomologia Experimentalis et Applicata*, **152**, 87-90.
- 645 van Langevelde F, Ettema JA, Donners M, Wallisdevries MF, Groenendijk D (2011) Effect of
 646 spectral composition of artificial light on the attraction of moths. *Biological Conservation*,
 647 **144**, 2274-2281.
- 648 VisitEngland (2012) The evening and night-time economy: realising the potential for destination
 649 organisations. VisitEngland & the Association of Town Centre Management.

- 650 Wakefield A, Stone EL, Jones G, Harris S (2015) Light-emitting diode street lights reduce last-ditch
651 evasive manoeuvres by moths to bat echolocation calls. *Royal Society Open Science*, **2**,
652 150291.
- 653 Warrant E, Dacke M (2016) Visual navigation in nocturnal insects. *Physiology*, **31**, 182-192.
- 654 West KE, Jablonski MR, Warfield B, Cecil KS, James M, Ayers MA,...Hanifin JP (2011) Blue light
655 from light-emitting diodes elicits a dose-dependent suppression of melatonin in humans.
656 *Journal of Applied Physiology*, **110**, 619-626.
- 657 Witherington BE (1992) Behavioral responses of nesting sea turtles to artificial lighting.
658 *Herpetologica*, **48**, 31-39.
- 659 Wollast R (1998) Evaluation and comparison of the global carbon cycle in the coastal zone and in the
660 open ocean, in *The Sea*, vol. 10, *The Global Coastal Ocean: Processes and Methods (Ideas*
661 *and Observations on the Progress of the Studies of the Seas)*, edited by K. H. Brink, and A. R.
662 Robinson, John Wiley, New York, pp. 213–252.
- 663 Wright KP, Mchill AW, Birks BR, Griffin BR, Rusterholz T, Chinoy ED (2013) Entrainment of the
664 human circadian clock to the natural light-dark cycle. *Current Biology*, **23**, 1554-1558.
- 665 Young CM, Chia F-S (1982) Ontogeny of phototaxis during larval development of the sedentary
666 polychaete, *Serpula vermicularis* (L.). *The Biological Bulletin*, **162**, 457-468.
- 667 Zissis G, Bertoldi P (2014) 2014 Update on the status of the LED market. European Commission
668 Report EU 27000 E
- 669 Zoltan K (2010) Measuring and modelling light pollution at the Zselic Starry Sky Park. *Journal of*
670 *Physics: Conference Series*, **218**, 012001.

Figure 1. The trend in research outputs associated with light pollution and climate change since the year 2000. Bar heights represent the cumulative number of articles expressed as a percentage of the total number of articles published by the end of 2016; numbers are the cumulative number of articles published by the end of each year. Note that the total number of articles does not reflect the total number published in the research area, only the number returned from the search. The data were collected from a Web of Science search for phrases in article titles. The search phrases used for light pollution research outputs were "Light pollution" OR "Artificial Light at Night" OR "Nighttime lighting" OR "Night-time lighting" OR "Night time lighting" OR "Street Lighting" OR "LED lighting" OR "Light-emitting diode lighting". The search phrase for climate change was 'Climate change' and results were not refined by research area. The search for articles on light pollution was refined by research areas: (PLANT SCIENCES OR ORNITHOLOGY OR PSYCHOLOGY MULTIDISCIPLINARY OR ENVIRONMENTAL SCIENCES OR EVOLUTIONARY BIOLOGY OR PHYSICS APPLIED OR ENTOMOLOGY OR ENGINEERING ENVIRONMENTAL OR ECOLOGY OR URBAN STUDIES OR FISHERIES OR BIODIVERSITY CONSERVATION OR BIOLOGY OR PHYSICS MULTIDISCIPLINARY OR ZOOLOGY OR OCEANOGRAPHY OR GEOGRAPHY PHYSICAL OR GEOGRAPHY OR REMOTE SENSING OR PHYSIOLOGY OR MARINE FRESHWATER BIOLOGY OR PUBLIC ENVIRONMENTAL OCCUPATIONAL HEALTH).


Figure 2. A comparison of fine scale spatial variability in environmental warming and artificial light at night on the Lizard peninsula, Cornwall, UK. A) The increase in the number of growing degree-days (a measure of a measure of change in growing season length expressed in °C Days) between 1977 and 2014 (100m resolution). Adapted with permission from Maclean *et al.* (2016). B) The distribution of artificial light across the same area (750m resolution) recorded from the VIIRS sensor on board the Suomi NPP satellite.

Figure 3. The potential ecological impacts of white Light Emitting Diode lighting compared to other light sources. Spectral power distributions are given for white Light Emitting Diode (LED),


Low Pressure Sodium (LPS), High Pressure Sodium (HPS) and Metal Halide (MH) lights recorded using a MAYA 200 pro spectrometer from street lighting in Cornwall. The amount of light at each wavelength is standardised to relative intensity (radiant energy divided by the maximum radiant energy recorded at any wavelength for each light source) so that the relative distribution of radiant energy across the light spectrum can be compared for each light source. Grey arrows represent the wavelength range over which different types of biological response are expected/recorded. Dashed lines represent the range of wavelengths over which Mammal, Bird, Reptile, Insect, and Arachnid visual systems can detect light [adapted from Davies *et al.* (2013)].

Figure 4. The sensitivity of marine invertebrates to direct artificial light and artificial sky glow.


Solid lines represent the attenuation of scalar irradiance (between 400 and 700nm) with depth estimated using radiative transfer models under winter (**a & c**; Chlorophyll = 0.3 mg m^{-3} uniform profile, wind = 5 m s^{-1}) and spring (**b & d**; Chlorophyll = 5 mg m^{-3} uniform profile, wind = 5 m s^{-1}) water column properties. Models of scalar irradiance with depth are derived from spectral power distribution recorded from the spring high tide mark under a white LED street light on the Barbican in Plymouth (**a & b**), and artificial skyglow from predominantly white Metal Halide spectra recorded above Falmouth Harbour (**c & d**). Grey dashed lines indicate the maximum depth at which sufficient artificial light is available to perform species behaviours. SSS= Settlement Site Selection; PR=Polyp Retraction; LP=Larval Phototaxis; DVM=Diel Vertical Migration. Sensitivities to white light were calculated from experimentally derived values in existing literature (Crisp & Ritz, 1973, Young & Chia, 1982, Forward *et al.*, 1984, Svane & Dolmer, 1995, Tankersley *et al.*, 1995, Båtnes *et al.*, 2013, Gorbunov & Falkowski, 2002).




159x69mm (300 x 300 DPI)



80x37mm (300 x 300 DPI)



150x99mm (300 x 300 DPI)



179x231mm (300 x 300 DPI)